



## Background

Recently, native and managed colonies of pollinators, including bees, butterflies, and other insects, have experienced a large-scale declines. In an effort to restore habitat for pollinators, the United States Department of Agriculture's Farm Service Agency (FSA) created the CP-42 Pollinator Habitat conservation practice under the Conservation Reserve Program. This program pays private landowners to restore habitat for pollinator species with ecological and economic importance. To qualify for the program, CP-42 seed mixes must include:

- A minimum of 9 pollinator-friendly flowering plants,
- A minimum of three species in bloom during each of three seasonal periods (April-June 15, June 15-July, August-October), and
- no more than 25% native, non-sod forming grasses.

Since 2014, more than 200,000 acres have been enrolled in the CP-42 program in Iowa with virtually no post-seeding monitoring or evaluation. To determine whether CP-42 sites are likely to meet the program's floral resource targets, we analyzed a 5-year data set on flower phenology from an existing prairie restoration established with a similar seed mix.

## Methods

- Data come from a long-term field experiment in prairie restoration at the Cedar River Ecological Research Site in Black Hawk County, Iowa. See Myers et al (2015) for details on site establishment and management.
- We analyzed 5 years of flower phenology data from a 32 species seed mix (P32) that included 20 forb and 12 graminoid species.
- The P32 forb seeding rate (26 pure live seeds/sq ft) was comparable to a typical CP-42 seed mix (~30 forb seeds/sq ft); however, the graminoid component greatly exceeded (55 pls/sq ft) CP-42 guidelines (~10 grass seeds/sq ft). However, other studies have illustrated that diverse mixes with higher grass to forb seeding ratios perform as well or better than forb-heavy pollinator mixes in producing nectar plants (Meissen et al., 2017)

## Research Questions


- Are there three P32 species in bloom during each of the CP-42 periods? Which species are most abundant?
- Does P32 habitat quality remain suitably high over 5 years? Is there an increase or decrease in flower abundance over time?

Table 1. P32 seed mix composition.

Scientific name*	Common name	Functional group
<i>Panicum virgatum</i>	switchgrass	C4 grass
<i>Andropogon gerardii</i>	big bluestem	C4 grass
<i>Schizachyrium scoparium</i>	little bluestem	C4 grass
<i>Bouteloua curtipendula</i>	side-oats grama	C4 grass
<i>Sorghastrum nutans</i>	indiangrass	C4 grass
<i>Agropyron smithii</i>	western wheatgrass	C3 grass
<i>Elymus canadensis</i>	Canada wildrye	C3 grass
<i>Elymus virginicus</i>	Virginia wildrye	C3 grass
<i>Astragalus canadensis</i>	milk vetch	Legume
<i>Desmodium canadense</i>	showy tick trefoil	Legume
<i>Heliopsis helianthoides</i>	ox-eye sunflower	Forb
<i>Lespedeza capitata</i>	round-headed bush clover	Legume
<i>Oligoneuron rigidum</i>	stiff goldenrod	Forb
<i>Ratibida pinnata</i>	yellow coneflower	Forb
<i>Helianthus grosseserratus</i>	saw-tooth sunflower	Forb
<i>Silphium laciniatum</i>	compass plant	Forb
<i>Carex bicknellii</i>	copper-shoulder oval sedge	Sedge
<i>Carex brevior</i>	plains oval sedge	Sedge
<i>Carex grvida</i>	long-awned bracted sedge	Sedge
<i>Sporobolus compositus</i>	composite dropseed	C4 grass
<i>Amorpha canescens</i>	leadplant	Legume
<i>Artemisia ludoviciana</i>	prairie sage	Forb
<i>Dalea purpurea</i>	purple prairie clover	Legume
<i>Echinacea pallida</i>	pale purple coneflower	Forb
<i>Eryngium yuccifolium</i>	rattlesnake master	Forb
<i>Monarda fistulosa</i>	wild bergamot	Forb
<i>Symphotrichum laeve</i>	smooth blue aster	Forb
<i>Symphotrichum novae-angliae</i>	New England aster	Forb
<i>Tradescantia bracteata</i>	longbract spiderwort	Forb
<i>Zizia aurea</i>	golden alexanders	Forb
<i>Phlox pilosa</i>	downy phlox	Forb
<i>Baptisia leucantha</i>	white wild indigo	Legume

## Results

Figure 1. Total mean abundance of sown flowers, weeds, and all flowers in P32 plots during 5 seasonal periods (1 = June 1 to June 20; 2 = June 21 to July 10; 3 = July 11 to July 31; 4 = August 1 to 31; 5 = Sept 1 to Sept 31) in a) 2010, b) 2011, c) 2012, d) 2013, and e) 2014



### Potential Management & Environmental Effects:

- 2010: establishment year / no management
- 2011: spring prescribed burn
- 2012: spring biomass harvest (summer drought)
- 2013: no management (May/June floods)
- 2014: spring prescribed burn (June floods)

### Photo of a P32 prairie planting



Figure 2. P32 survey sessions 1-5 re-formatted to fit CP-42 bloom periods for: a) sown species, b) weeds, and c) total flowers. Dashed red line indicates a bloom threshold of 2 flowers per m<sup>2</sup>.




Table 1. Total # of species in bloom during CP-42 periods at various bloom abundance thresholds

Year	Total # sown species	PERIOD 1 (April to June 15)		
		Sown > 1 flower per 10 m <sup>2</sup>	Sown > 1 flower per m <sup>2</sup>	Weeds (>1 flower per m <sup>2</sup> )
2010	6	4	2 (hehe, phpi)	9 (1, meof)
2011	4	3	1 (ziau)	1 (0)
2012	5	2	0	3 (0)
2013	3	2	0	2 (0)
2014	4	2	1 (trbr)	1 (0)

Year	Total # sown species	PERIOD 2 (June 15 to July)		
		Sown > 1 flower per 10 m <sup>2</sup>	Sown > 1 flower per m <sup>2</sup>	Weeds (>1 flower per m <sup>2</sup> )
2010	9	6	1 (hehe)	11 (1, meof)
2011	11	8	1 (hehe)	3 (0)
2012	14	4	1 (hehe)	3 (0)
2013	9	3	0	3 (0)
2014	11	3	0	0

Year	Total # sown species	PERIOD 3 (August to October)		
		Sown > 1 flower per 10 m <sup>2</sup>	Sown > 1 flower per m <sup>2</sup>	Weeds (>1 flower per m <sup>2</sup> )
2010	11	7	2 (deca, hehe)	14 (0)
2011	12	9	2 (deca, hehe)	5 (0)
2012	10	4	1 (hehe)	4 (0)
2013	12	5	1 (deca)	3 (0)
2014	14	7	1 (deca)	3 (1, sypl)

## Conclusions

- ❖ Floral resources were dominated by sown species, and flower abundance declined greatly after Year 2.
- ❖ Total and sown flower abundance was < 2 flowers per m<sup>2</sup> during Period 1 in 2012 and 2013 and during Period 2 in 2013 and 2014.
- ❖ The most abundant sown species were *Phlox pilosa*, *Zizia aurea*, and *Tradescantia bracteata* in Period 1, *Heliopsis helianthoides* in Period 2, and *Heliopsis helianthoides* and *Desmodium canadense* in Period 3.
- ❖ After Year 1, weeds contributed only a minor portion of total flower abundance. Early season sweet clovers and late season frost aster and goldenrods were the most common weeds.
- ❖ With no density threshold, the P32 seed mix met the requirement of 3 species in bloom during each CP-42 seasonal period.
- ❖ With a bloom threshold of 1 flower per 10 m<sup>2</sup>, the site did not meet the requirement of 3 species in bloom during Period 1 in 2012, 2013, or 2014.
- ❖ With a bloom threshold of 1 flower per 1 m<sup>2</sup>, the site did not meet the requirement of 3 species in bloom during any seasonal period in any year.
- ❖ While the P32 seed mix met the general requirements of the CP-42 program, the abundance and diversity of floral resources declined over time, particularly in the first seasonal period (April-June 15).

## Acknowledgements

I would like to thank the Tallgrass Prairie Center for allowing me to contribute to this research project, Mark Myers for all of his time and support, and the Carver Foundation for the funding that made this poster possible.

### References

Myers, M.C., J.T. Mason, B.J. Hoksch, C.A. Cambardella, J.D. Pfrimmer. 2015. Birds and butterflies respond to soil-induced habitat heterogeneity in experimental plantings of tallgrass prairie species managed as agroenergy crops in Iowa, USA. *Journal of Applied Ecology* 52(5):1176-1187. DOI: 10.1111/1365-2664.12503

Meissen, Justin; Williams, Dave; and Jackson, Laura. 2017. Cost Effective Native Seed Mix Design and First-Year Management. *Farm Progress Reports* 2016(1): Article 62. Available from: <https://lib.dr.iastate.edu/farmprogressreports/vol2016/iss1/62>

United States Department of Agriculture, Farm Service Agency. 2011. *Practice CP42, Pollinator Habitat* (Notice CRP-687). Washington D.C.

United States Department of Agriculture, Natural Resources Conservation Service. 2015. PLANTS Database. Available from: <http://plants.usda.gov/java/>.